

NEW YORK STATE COLLEGE OF AGRICULTURE
CORNELL UNIVERSITY
ITHACA, NEW YORK
LABORATORY OF BACTERIOLOGY

April 26, 1949

Dr. Joshua Lederberg
Department of Genetics
University of Wisconsin
Madison 6, Wisconsin

Dear Josh:

Enclosed are a couple more pedigrees from H168. I was a little suspicious of the small one so did not want to invest the time necessary to carry it any farther. The original cell proved to be a heterozygote so it would have been all right. The series 5 cultures, I think, are the most extensive group yet and I intend to carry ^{all of} them out this far, ^{in the future} hoping that the segregation will occur early enough that we can perhaps account for the possible inviable nucleus. This series 5 case, cultures 239 and 240 with the inviable 5-120, would be interesting if we did not already have several observations indicating that the sib cell to a segregant is a viable heterozygote.

I am also sending out the following cultures from the 5 series:

203, 204, 195, 196, 25, 53, 54

just on the chance that they may be valuable, although I think probably all of them are heterozygotes. I am checking them here further but I think it is best to get them to you as quickly as possible. I am dumping some of the original broth culture into some melted agar to attempt to cut down the amount of growth before the cultures get to you.

I am not planning on giving anything at Cincinnati. If you wish to use any of the stuff we have so far you are welcome to do so. Actually, there is not a great deal to say beyond the indication that the mechanism is more complex than one might have hoped for. I hope to have several more pedigrees as extensive as this series 5 before the Cincinnati meetings.

Have you played around with cold shocks or any other mechanism for increasing the frequency of the segregation? It would be a big help to me if there were some way of inducing the segregation to occur at a bit higher rate. I have made a few random notes on the back of the series 5 sheets which may or may not indicate that I can spot a cell which will not grow. Actually, I think if I were more careful I could raise my predicting ability to something like 80 per cent accuracy. The main reason I am interested in this is that I feel the failure of cells to grow can not be laid entirely to the rigors of the technique and that there is an actual physiological difference which arises before I separate the sibs. As to culture 2-200 I am positive that at

HP

April 26, 1949

the time I plated ~~it~~ truly segregating heterozygous cells were present in the culture. I will replate in an effort to confirm this. Actually I have never classified a culture as heterozygous on the basis of its being a mixture of + and - cells. I raised the question earlier because it was a possibility.

What is your guess as to the nature of the aberrant region? Just ~~that~~ ^{plus} a deficiency or a deficiency ~~shows~~ ^{after} some to the change or is the situation so confused that one can not make a good guess?

Does ~~Mt1-~~ revert readily? If so, I will switch to xylose.

Very truly yours,

M. R. Zelle
M. R. Zelle

MRZ:jc

Enclosure

fresh,
#72 was derived from a cross of
B M Het x T L B₁ loc-Mal-Gal-Hyl-Arab-V₁^r
but was + for Mal, Gal, Arab. Is this correct?

Also - #168 is homo or hemi-zygous for Mal-, arab-, ???

Chamber for the xylose, our came in the same day I got your letter.

H168 genotype? = $\frac{B \ M \ V_k^+ \ Het \ + \ + \ + \ + \ + \ + \ + \ +}{+ \ + \ + \ + \ T \ L \ B_1 \ loc, \ Mal \ Hyl \ Gal \ Arab \ Malt \ V_1^+}$